

FIRST AID FACT SHEET

Blue Ringed Octopus and Cone Shell Bites

Blue Ringed Octopus are commonly found in tidal pools, and if handled may inflict a potentially fatal bite. This may not be painful, as venom is not always injected (envenomation).

The Cone Shell is mainly found in tropical waters, envenomation happens when they are handled or stepped on and they fire a dart like barb which delivers venom.

Although different, venoms from both these creatures can cause paralysis and death from respiratory failure within 30 minutes.

What to do

1. Follow DRSABCD St John Action Plan.
2. Calm casualty and keep them still.
3. If bite is on a limb apply a firm broad crepe or elasticised roller bandage starting just above the fingers or toes, and moving upwards on the bitten limb as far as can be reached and immobilise the limb with a splint.
4. Monitor casualty closely for signs of adverse reactions or deterioration.
5. Urgent medical aid. Call **Triple Zero (000)** for an ambulance.
6. Prepare to perform cardiopulmonary resuscitation (CPR) if necessary.

Caused by

- + Handling the creature.
- + Accidentally standing on the creature.

Signs & symptoms

- + A painless bite with a spot of blood visible.
- + Numbness to lips and tongue, muscle weakness which can stop the casualties breathing.



In a medical emergency
call **Triple Zero (000)**

Book a St John first aid course today | (08) 9334 1233 | [stjohnambulance.com.au](https://www.stjohnambulance.com.au)

Information provided is as per Australian Resuscitation Council Guidelines. These first aid protocols are for the Australian market only. All care has been taken in preparing this information but St John takes no responsibility for its use. This information is not a substitute for practical first aid training with St John.