

## FIRST AID FACT SHEET

# Red-Back Spider Bite

A Red-Back spider bite can be life threatening to a child, but apart from the pain it is rarely serious for an adult.

They are common in dry places around buildings, outdoor furniture, machinery and stacked materials. In the bush, they nest under logs and rocks.

### Caused by

- + Spider biting and puncturing the skin.

### Signs & symptoms

- + Immediate pain at the bite site which becomes hot, red and swollen.
- + Intense local pain which increases and spreads.
- + Nausea, vomiting and abdominal pain.
- + Profuse sweating, especially at the bite site.
- + Swelling of glands in the groin or armpit of the limb that was bitten.

### What to do

1. Follow DRSABCD St John Action Plan.
2. Lay the casualty down, rest and reassure.
3. Monitor the casualty constantly.
4. Apply a cold compress/cold pack to lessen the pain (no longer than twenty (20) minutes).
5. Seek medical aid promptly, if:
  - + Person bitten is a young child, elderly or infirmed.
  - + The casualty collapses.
  - + Pain is severe.



In a medical emergency  
call **Triple Zero (000)**

Book a St John first aid course today | (08) 9334 1233 | [stjohnambulance.com.au](https://stjohnambulance.com.au)

Information provided is as per Australian Resuscitation Council Guidelines. These first aid protocols are for the Australian market only. All care has been taken in preparing this information but St John takes no responsibility for its use. This information is not a substitute for practical first aid training with St John.